

**Combining environmental suitability and population abundances to evaluate the invasive potential of the tunicate *Ciona intestinalis* along the temperate South American coast**

Stella Januario<sup>1\*</sup>; Sergio A. Estay<sup>2,4</sup>, Fabio A. Labra<sup>3</sup>, Mauricio Lima<sup>1,4</sup>

<sup>1</sup> *Departamento de Ecología, Pontificia Universidad Católica de Chile, Alameda 340, Santiago, Chile*


<sup>2</sup> *Instituto de Ciencias Ambientales y Evolutivas, Facultad de Ciencias. Universidad Austral de Chile. Casilla 567, Valdivia, Chile*

<sup>3</sup> *Centro de Investigación e Innovación para el Cambio Climático, Facultad de Ciencias, Universidad Santo Tomás, Ejercito 146, Código Postal 8370003, Santiago, Chile.*

<sup>4</sup> *Center of Applied Ecology & Sustainability (CAPES), Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, Santiago 6513677, Chile.*

## Supplementary figures

**Figure 1.** Projections of the potential distribution of *C. intestinalis* on Chilean coast using non analog environments for each of the five locations used in the analysis.



**Figure 2.** Projections of the potential distribution of *C. intestinalis* on Chilean coast using just analog environments for each of the five locations used in the analysis.

**Non-analog environments**

